arXiv:1510.05974v1 [math.FA] 20 Oct 2015

Distortion in the finite determination result for embeddings of locally finite metric spaces into Banach spaces

Sofiya Ostrovska and Mikhail I. Ostrovskii

Abstract

Given a Banach space X and a locally finite metric space A, it is known that if all finite subsets of A admit bilipschitz embeddings into X with distortions $\leq C$, then the space A itself admits an embedding into X with distortion $\leq D \cdot C$, where D is an absolute constant. The goal of this paper is to show that $D > 1$, implying that, in general, there is a "deterioration of distortion" in the aforementioned situations.

Keywords. Banach space, distortion of a bilipschitz embedding, locally finite metric space

2010 Mathematics Subject Classification. Primary: 46B85; Secondary: 46B20.

1 Introduction

The study of bilipschitz embeddings of metric spaces into Banach spaces is a very active research area which has found many applications, see [3, 4, 6]. In this paper, we contribute to the study of relations between the embeddability of an infinite metric space and its finite pieces. Let us recollect some necessary notions.

Definition 1.1. A metric space is called *locally finite* if each ball of finite radius in it has finite cardinality.

Definition 1.2. (i) Let $0 \leq C < \infty$. A map, say $f : (A, d_A) \to (Y, d_Y)$, between two metric spaces is called C-*Lipschitz* if

$$\forall u, v \in A \quad d_Y(f(u), f(v)) \leq C d_A(u, v).$$

A map f is called *Lipschitz* if it is C-Lipschitz for some $0 \leq C < \infty$.

(ii) Let $1 \leq C < \infty$. A map, $f : A \to Y$, is called a C-*bilipschitz embedding* if there exists $r > 0$ such that

$$\forall u, v \in A \quad r d_A(u, v) \leq d_Y(f(u), f(v)) \leq r C d_A(u, v). \tag{1}$$

A *bilipschitz embedding* is one which is C-bilipschitz for some $1 \leq C < \infty$. The smallest constant C for which there exists $r > 0$ such that (1) is satisfied, is called the *distortion* of f.

It has been known that the bilipschitz embeddability of locally finite metric spaces into Banach spaces is finitely determined in the following sense:

Theorem 1.3 ([5]). *Let A be a locally finite metric space whose finite subsets admit bilipschitz embeddings with uniformly bounded distortions into a Banach space X. Then, A also admits a bilipschitz embedding into X.*

To elaborate more, the argument of [5] implies that there exists an absolute constant D, such that, if all finite subsets of A admit bilipschitz embeddings into X with distortions $\leq C$, then A admits an embedding into X with distortion $\leq D \cdot C$. It is well known that if a Banach space X is such that each separable subspace of any ultrapower of X admits an isometric embedding into X, then the statement of the previous paragraph holds with $D = 1$.

The main goal of this paper is to show that, for general Banach spaces X, the constant D is strictly greater than 1.

Remark 1.4. The result of [5] was preceded with one established by Baudier and Lancien [1], who proved that each locally finite metric space admits a bilipschitz embedding into any Banach space admitting bilipschitz embeddings of $\{\ell_\infty^n\}_{n=1}^\infty$ with uniformly bounded distortions. Since space X in our construction is of the form $X = (\oplus_{n=1}^\infty \ell_\infty^n)_2$, it follows that the "deterioration of distortion" also occurs in this case.

2 The main result

The following result immediately shows that for general Banach spaces, D is not equal to 1 and, thence, $D > 1$.

Theorem 2.1. *There exists a locally finite metric space A and a Banach space X such that the finite subsets of A admit isometric embeddings into X while A itself does not admit an isometric embedding into X.*

Proof. Consider $X = (\oplus_{n=1}^\infty \ell_\infty^n)_2$. We refer to [6, Definition 3.6] for the definition of such direct sums. For this X, the condition pertaining to the finite subsets is satisfied for all locally finite metric spaces A because, by the result of Fréchet [2, p. 161] (see also [6, Proposition 1.17]), each n-element metric space admits an isometric embedding into ℓ_∞^n. Therefore, what remains is to build A in such a way that it will not be isometric to a subset of X.

To begin with, consider triples of points $x, y, z \in X$ satisfying

$$||x - z|| = ||x - y|| + ||y - z||. \tag{2}$$

Let $x = \{x_n\}$, $y = \{y_n\}$, $z = \{z_n\}$, where $x_n, y_n, z_n \in \ell_\infty^n$ are the components of x, y, and z, respectively. Now comes the following simple observation.

Lemma 2.2. *For any triple $x, y, z \in X$ satisfying* (2)*, the vector $\{||x_n - y_n||\}_{n=1}^\infty \in \ell_2$ is a positive multiple of $\{||y_n - z_n||\}_{n=1}^\infty \in \ell_2$.*

Proof. Assuming the contrary, one derives that the ℓ_2-norm of the vector $\{||x_n - y_n|| + ||y_n - z_n||\}_{n=1}^{\infty}$ is strictly less than

$$\|\{||x_n - y_n||\}_{n=1}^{\infty}\|_2 + ||\{||y_n - z_n||\}_{n=1}^{\infty}||_2 = ||x - y||_X + ||y - z||_X.$$

On the other hand,

$$\|\{||x_n - y_n|| + ||y_n - z_n||\}_{n=1}^{\infty}\|_2 \geq \|\{||x_n - z_n||\}_{n=1}^{\infty}\|_2 = ||x - z||_X,$$

which contradicts (2). □

Definition 2.3. A *metric ray* in a metric space (A, d_A) is a sequence $R = \{r_i\}_{i=0}^{\infty}$ of points such that the sequence $d_A(r_i, r_0)$ is strictly increasing and, for $i < j < k$, the following equality holds:

$$d_A(r_i, r_k) = d_A(r_i, r_j) + d_A(r_j, r_k). \tag{3}$$

For all of the metric rays in the Banach spaces considered in this paper, it will be assumed that

$$r_0 = 0. \tag{4}$$

Consider subspaces $X_k = \left(\oplus_{n=1}^{k} \ell_\infty^n\right)_2$ in X and the natural projections $P_k : X \to X_k$ defined by $P(\{x_n\}_{n=1}^{\infty}) = \{x_n\}_{n=1}^{k}$.

Lemma 2.4. *For each metric ray $R = \{r_i\}_{i=0}^{\infty}$ in X and each $\varepsilon \in (0, 1)$, there is $k \in \mathbb{N}$ such that the natural projection $P_k : X \to X_k$ satisfies:*

$$||P_k r_i - r_i|| \leq \varepsilon ||r_i|| \tag{5}$$

Furthermore, a number k satisfying this condition can be determined using only the number $\varepsilon > 0$ and the vector r_1, as $r_0 = 0$ by (4).

Proof. Let $r_i = \{r_{in}\}_{n=1}^{\infty}$, where $r_{in} \in \ell_\infty^n$. Combining Lemma 2.2 and the definition of a metric ray, we conclude that for $i < j < k$, the vector $\{||r_{jn} - r_{in}||\}_{n=1}^{\infty}$ is a positive multiple of $\{||r_{kn} - r_{jn}||\}_{n=1}^{\infty}$. Using the fact that $r_{0n} = 0$ for every n, it can be easily obtained that any vector of the form $\{||r_{jn} - r_{in}||\}_{n=1}^{\infty}$ is a positive multiple of $\{||r_{1n}||\}_{n=1}^{\infty}$. Now, pick $k \in \mathbb{N}$ such that $||P_k r_1 - r_1|| \leq \varepsilon ||r_1||$. The statement follows. □

Let the metric space A be a locally finite subset of ℓ_∞ formed as a union of a sequence $\{r(j)\}_{j=1}^{\infty}$ of metric rays $r(j) = \{r_i(j)\}_{i=0}^{\infty}$ satisfying the conditions:

- $r_0(j) = 0$ for all j.
- $r_1(j) \neq 0$ is the same for all j.

Suppose that A admits an isometric embedding $E : A \to X$. Clearly, isometries map metric rays onto metric rays. Without loss of generality, assume that $E(0) = 0$.

Hence, by Lemma 2.4, for each $\varepsilon \in (0,1)$, there is $k \in \mathbb{N}$ such that, for every ray $r(j)$ in A and $i \in \mathbb{N}$, one has:

$$||P_k E r_i(j) - E r_i(j)|| \leq \varepsilon ||E r_i(j)|| = \varepsilon ||r_i(j)||, \tag{6}$$

where the equality holds because E is an isometry mapping 0 to 0. It can be readily seen that k depends only on the vector Er_1. Notice that there is no control on the value of k and, thence on the dimension $m = \dim X_k$. Yet, to complete our argument, it suffices to know that for each A of the described form and each isometric embedding $E : A \to X$, this dimension is finite.

For the remainder of this paper, $\varepsilon \in (0,1)$ will be selected in such a way that, for every $t \in \mathbb{N}$,

$$3^t - 2\varepsilon 3^{t+1} \geq 3^{t-1}. \tag{7}$$

It is common knowledge that there exists an absolute constant C such that, for any $\delta > 0$, the cardinality of a δ-separated set in a ball of radius R in X_k with $\dim X_k = m$ does not exceed $(CR/\delta)^m$.

Therefore, to prove Theorem 2.1 it is enough to find a locally finite metric space A consisting of a union of metric rays as described above and in a way that the cardinalities of 3^{t-1}-separated sets in $P_k E B_t$, where B_t is a ball of radius 3^{t+1} centered at 0 of A, grow at least as fast as an a-priori chosen sequence $\{N_t\}_{t=1}^{\infty}$.

Such a metric space A is constructed inside the countable subset S of ℓ_∞ consisting of all sequences, for which the first coordinate is a positive integer, the next N_1 coordinates are positive integer multiples of 3, the next N_2 coordinates are positive integer multiples of 3^2, the next N_3 coordinates are positive integer multiples of 3^3, and so on. With this in mind, it is not difficult to see that S is locally finite for an arbitrary sequence $\{N_k\}_{k=1}^{\infty}$ of finite numbers, and thus A is also locally finite.

Now, let us introduce A as a union of metric rays $r(j)$. For each $j \in \mathbb{N}$, pick one of the coordinates $n_1(j)$ of the N_1-bunch, one - $n_2(j)$ - of the N_2-bunch, etc. This will be performed in such a way that the following condition is satisfied:

$$\forall n \in N_k\text{-bunch } \exists j \in \mathbb{N} \quad n = n_k(j). \tag{8}$$

After this, the metric rays are defined as follows:

1. $r_0(j) = 0$,
2. $r_1(j)$ is the unit vector $(1, 0, \dots, 0, \dots) \in \ell_\infty$.
3. $r_2(j)$ has $1 + 3$ as the first coordinate, 3 as the $n_1(j)$ coordinate, and all the other coordinates are 0;
4. $r_3(j)$ has $1 + 3 + 3^2$ as the first coordinate, $3 + 3^2$ as the $n_1(j)$ coordinate, 3^2 as the $n_2(j)$ coordinate, and all the other coordinates are 0;
5. Similarly, for all $r_k(j)$, $k \geq 4$.

It can be noticed that each $r(j)$ is a metric ray. Also, for any $t \in \mathbb{N}$, the vector $r_{t+1}(j)$ is in B_t and even in the ball of radius $1 + 3 + 3^2 + \cdots + 3^t$. Combining the definition of $r(j)$ and the condition (8), it is concluded that the set of all vectors $r_{t+1}(j)$ contains a subset of cardinality N_t which is 3^t-separated.

Applying inequality (6) to any two elements $Er_{t+1}(j_1)$ and $Er_{t+1}(j_2)$ of this subset, what follows can be reached:

$$\begin{aligned} &||P_k Er_{t+1}(j_1) - P_k Er_{t+1}(j_2) - (Er_{t+1}(j_1) - Er_{t+1}(j_2))|| \\ &\qquad \leq ||P_k Er_{t+1}(j_1) - Er_{t+1}(j_1)|| + ||P_k Er_{t+1}(j_2) - Er_{t+1}(j_2)|| \\ &\qquad \leq \varepsilon(||r_{t+1}(j_1)|| + ||r_{t+1}(j_2)||) \end{aligned}$$

and, as a result,

$$||P_k Er_{t+1}(j_1) - P_k Er_{t+1}(j_2)|| \geq 3^t - 2\varepsilon 3^{t+1} \overset{(7)}{\geq} 3^{t-1},$$

which leads to our ultimate conclusion. □

3 Final remark

We have left open the arguably very challenging problem on the precise estimate for the "deterioration of distortion" when passing from the embeddings of finite subsets of a locally finite metric spaces to an embedding of the whole locally finite metric space.

The second-named author gratefully acknowledges the support by the National Science Foundation DMS-1201269.

The authors express their sincere gratitude to Mr. P. Danesh for very useful suggestions.

References

[1] F. Baudier, G. Lancien, Embeddings of locally finite metric spaces into Banach spaces, *Proc. Amer. Math. Soc.*, **136** (2008), 1029–1033.

[2] M. Fréchet, Les dimensions d'un ensemble abstrait, *Math. Ann.*, **68** (1910), no. 2, 145–168.

[3] J. Matoušek, *Lectures on Discrete Geometry*, Springer-Verlag, New York, 2002.

[4] A. Naor, L_1 embeddings of the Heisenberg group and fast estimation of graph isoperimetry, in: *Proceedings of the International Congress of Mathematicians*, 2010, Hyderabad India, vol. **III**, 2011, pp. 1549–1575.

[5] M. I. Ostrovskii, Embeddability of locally finite metric spaces into Banach spaces is finitely determined, *Proc. Amer. Math. Soc.*, **140** (2012), 2721–2730.

[6] M. I. Ostrovskii, *Metric Embeddings: Bilipschitz and Coarse Embeddings into Banach Spaces*, de Gruyter Studies in Mathematics, **49**. Walter de Gruyter & Co., Berlin, 2013.

Department of Mathematics, Atilim University, 06836 Incek, Ankara, TURKEY
E-mail address: sofia.ostrovska@atilim.edu.tr

Department of Mathematics and Computer Science, St. John's University, 8000 Utopia Parkway, Queens, NY 11439, USA
E-mail address: ostrovsm@stjohns.edu